

**AD-A273 298**



(1)

OFFICE OF NAVAL RESEARCH

GRANT : N00014-89-J-3062

R&T Code 4132084

Technical Report No. 24

Synthesis of Polyphenylenes via Bergman Cyclization of Enediyne Monomers

by

Tour, J. M.; John, J. A.

Accepted for Publication in

Polym. Prepr. (Am. Chem. Soc., Div. Polym. Chem.)

**S** DTIC  
ELECTE  
DEC 01 1993  
**A**

Department of Chemistry and Biochemistry  
University of South Carolina  
Columbia, SC 29208

November 22, 1993

Reproduction in whole, or in part, is permitted for any purpose of the United States Government

This document has been approved for public release and sale; its distribution is unlimited.

**93-29301**



296

**93 11 30018**


REPORT DOCUMENTATION PAGE			Form Approved OMB No 0704-0188	
<small>Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188) Washington, DC 20503.</small>				
1. AGENCY USE ONLY (Leave blank)	2. REPORT DATE 11/22/93	3. REPORT TYPE AND DATES COVERED Technical Report		
4. TITLE AND SUBTITLE  Synthesis of Polyphenylenes via Bergman Cyclization of Eneidyne Monomers.		5. FUNDING NUMBERS  G- N00014-89-J3062 R&T 4132084		
6. AUTHOR(S)  Tour, J. M.; John, J. A.				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Department of Chemistry and Biochemistry University of South Carolina Columbia, SC 29208		8. PERFORMING ORGANIZATION REPORT NUMBER		
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) Department of the Navy Office of Naval Research 800 North Quincy Street Arlington, VA 22217-5000		10. SPONSORING / MONITORING AGENCY REPORT NUMBER  24		
11. SUPPLEMENTARY NOTES  <i>Polym. Prepr. (Am. Chem. Soc., Div. Polym. Chem.) 1993, 34(2), 372.</i>				
12a. DISTRIBUTION / AVAILABILITY STATEMENT Reproduction in whole or in part is permitted for any purpose of the United States Government. This document has been approved for public release and sale; its distribution is unlimited.			12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words)  Described is the synthesis of polyphenylenes via the Bergman cyclization of enediynes monomers. These monomers were synthesized via the Stephans-Castro/Sonogashira reaction and polymerized under varying conditions.				
14. SUBJECT TERMS			15. NUMBER OF PAGES	
			16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT UL	

## Synthesis of Polyphenylenes via Bergman Cyclization and Polymerization of Eneidyne Monomers

James M. Tour\* and Jens A. John  
Department of Chemistry and Biochemistry  
University of South Carolina  
Columbia, South Carolina 29208


In the early 1980s Bergman's study of thermal cycloaromatization of enediynes led to the suggestion of a 1,4-benzene diradical as an intermediate.<sup>1</sup> Later, DNA-cleaving antibiotics such as the esperamicins, calicheamicins and dynemicins were discovered to possess a cyclic enediyne moiety which underwent a Bergman cyclization upon activation.<sup>2</sup> Therefore synthetic and mechanistic studies were performed to investigate this reaction further by thermally cyclizing a wide variety of enediynes in the presence of 1,4-cyclohexadiene as a radical quencher.<sup>1,3</sup>

DTIC QUALITY INSPECTED 6




Since the natural products contained cyclic enediynes, most of the studies focused on these compounds. The model cyclizations in the presence of 1,4-cyclohexadiene resulted in moderate yields of substituted benzenes and polymeric by-products.

Our interest in the area was kindled when we regarded the intermediate 1,4-benzene diradical as a building block for substituted polyphenylenes.



Accession For		
NTIS	CRA&I	<input checked="" type="checkbox"/>
DTIC	TAB	<input type="checkbox"/>
Unannounced		<input type="checkbox"/>
Justification		
By		
Distribution /		
Availability Codes		
Dist	Avail and/or Special	
A-1		

In addition, benzene analogs of enediynes were envisioned to form 1,4-naphthalene diradical intermediates which in turn could lead to naphthalene-based polymers.




The targeted monomers all contained non-cyclic enediynes to ensure better stability and ease of synthesis.


Recently, Swager *et al.* used cyclic benzene analogs of enediynes to construct an all carbon ladder polymer.<sup>4</sup>

### Synthesis of Enediyne Monomers

Both alkenyl and aryl diyne monomers were synthesized via the Stephens-Castro/Sonogashira reaction<sup>5</sup> in which a vinyl or aryl halide was coupled with a terminal acetylene.



This provided rapid access to the required mono- and disubstituted alkynyl and aryl diyne monomers. A list of the monomers prepared is shown below.



These monomers were then polymerized under various conditions (Table I).

Monomer	Method <sup>a</sup> / Temp.	Table I.			solubility
		M <sub>w</sub>	M <sub>n</sub>	$\frac{M_w}{M_n}$	
1	A/170	-	-	-	insoluble
2	A/400	245	141	1.7	0.33 sol. 0.67 insol.
3	A/340	106	71	1.5	0.5 sol. 0.5 insol.
4	A/300	2850	1040	2.7	soluble
5	A/300	590	430	1.4	soluble

4	B/120	325	312	1.0	soluble
				4	
5	B/140	1880	790	2.4	soluble
6	B/120	3930	1570	2.5	soluble
5	D/150	915	522	1.7	soluble
7	B/150	305000	9500	32	soluble
7	C/140	2000	630	3.2	soluble
7	A/150	2000	630	3.2	soluble
7	E/47 to 118	3200	940	3.4	soluble
8	B/ 95	10600	4100	2.6	soluble

<sup>a</sup>Method A: heated neat. Method B: heated in benzene. Method C: added slowly to refluxing 1,2-dichlorobenzene. Method D: heated together with a radical initiator. Method E: slowly increased temperature over several hours.

### Discussion

These results led to several conclusions. First, the use of benzenoid monomers was not efficacious due to the lack of polymer solubility, low molecular weights and high polymerization temperatures required. Secondly, alkyl substituents in the monomer led to low molecular weights presumably due to radical transfer generating more stable alkyl, allylic, or benzylic radicals and ultimately causing premature chain termination. Last, substitution on both alkynes required higher polymerization temperatures. At these temperatures, decomposition of the monomer may compete with polymer growth. The best results were obtained when one of the alkynes was unsubstituted and the other one was bearing an aromatic group.

### Conclusion

This study has shown the feasibility of the polymerization of enediyne monomers via a Bergman cyclization to afford low molecular weight polyaromatics. Our results show that soluble polyphenylenes can be easily made from non-cyclic enediynes. Further studies with other aryl substituents are planned to narrow the polydispersity and increase the molecular weight.

**Acknowledgments.** We are grateful for support from the Office of Naval Research and the National Science Foundation (EHR-91-08772, DMR-9158315), and generous industrial contributors to the NSF Presidential Young Investigator Award (1991-96): Hercules, IBM, Ethyl, Shell, and Eli Lilly Corporations. We also thank Molecular Design Ltd. for the use of their synthetic data base.

(1) Lockhardt, T. P.; Comita, P. B.; Bergman, R. G. *J. Am. Chem. Soc.* 1981, 103, 4082. Lockhardt, T. P.; Bergman, R. G. *J. Am. Chem. Soc.* 1981, 103, 4091.

(2) Lee, M. D.; Dunne, T. S.; Siegel, M. M.; Chang, C. C.; Morton, G. O.; Borders, D. B. *J. Am. Chem. Soc.* 1987, 109, 3464. Lee, M. D.; Dunne, T. S.; Chang, C. C.; Ellestad, G. A.; Siegel, M. M.; Morton, G. O.; McGahren, W. J.; Borders, D. B. *J. Am. Chem. Soc.* 1987, 109, 3466. Golik, J.; Clardy, J.; Dubay, G.; Groenewold, G.; Kawaguchi, H.; Konishi, M.; Krishnan, B.; Ohkuma, H.; Saitoh, K.; Doyle, T. W. *J. Am. Chem. Soc.* 1987, 109, 3461. Golik, J.; Dubay, G.; Groenewold, G.; Kawaguchi, H.; Konishi, M.; Krishnan, B.; Ohkuma, H.; Saitoh, K.; Doyle, T. W. *J. Am. Chem. Soc.* 1987, 109, 3462. Konishi, M.; Ohkuma, H.; Matsumoto, K.; Tsuno, T.; Kamei, H.; Miyaki, T.; Oki, T.; Kawaguchi, H.; VanDuyne, G.D.; Clardy, J. *J. Antibiot.* 1989, 42, 1449. For a recent review see: Nicolaou, K. C.; Dai, W.-M. *Angew. Chem. Int. Ed. Engl.* 1991, 30, 1387.

(3) Beau, J. M.; Crevisy, C. *Tetrahedron Lett.* 1991, 32, 3171. Singh, R.; Just, G. *Tetrahedron Lett.* 1991, 32, 185. Nagata, R.; Yamanaka, H.; Murahashi, E.; Saito, I. *Tetrahedron Lett.* 1990, 31, 2907. Nicolaou, K.C.; Maligres, P.; Shin, J.; de Leon, E.; Rideout, D. *J. Am. Chem. Soc.* 1990, 112, 7825. Wender, P. A.; McKinney, J. A.; Mukai, C. *J. Am. Chem. Soc.* 1990, 112, 5369.

(4) Zhou, Q.; Swager, T. M. *Polym. Prepr. (Am. Chem. Soc., Div. Polym. Chem.)* 1993, 34, 193.

(5) Sonogashira, K.; Tohda, Y.; Hagihara, N.; *Tetrahedron Lett.* 1975, 4467. Stephans, R. D.; Castro, C. E. *J. Org. Chem.* 1963, 28, 3313. Suffert, J.; Ziesel, R. *Tetrahedron Lett.* 1991, 32, 757.